arXiv:math-ph/0310046v1 23 Oct 2003

Jets of modules in noncommutative geometry

G. Sardanashvily

Department of Theoretical Physics, Moscow State University, 117234 Moscow, Russia
E-mail: sard@grav.phys.msu.su
URL: http://webcenter.ru/∼sardan/

Abstract. Jets of modules over a commutative ring are well known to make up the representative objects of linear differential operators on these modules. In noncommutative geometry, jets of modules provide the representative objects only of a certain class of first order differential operators. As a consequence, a generalization of the standard Lagrangian formalism on smooth manifolds to noncommutative spaces is problematic.

Let $\mathcal{K}$ be a commutative ring and $\mathcal{A}$ a commutative $\mathcal{K}$-ring. Let P and Q be $\mathcal{A}$-modules and $\mathcal{J}^r(P)$ the module of r-order jets of P. There is a $\mathcal{A}$-module isomorphism

$$\mathrm{Hom}_{\mathcal{A}}(\mathcal{J}^r(P), Q) = \mathrm{Diff}_r(P, Q), \tag{1}$$

where $\mathrm{Diff}_r(P, Q)$ is the module of r-order Q-valued linear differential operators on an $\mathcal{A}$-module P [4]. It follows that the jet module $\mathcal{J}^r(P)$ is the representative object of the functor $Q \to \mathrm{Diff}_r(P, Q)$.

In particular, let $\mathcal{A} = C^\infty(X)$ be the $\mathbb{R}$-ring of smooth real functions on a smooth manifold X. If P is a projective $C^\infty(X)$-module of finite rank, it is isomorphic to the module $Y(X)$ of global sections of some vector bundle $Y \to X$. In this case, $\mathcal{J}^r(P)$ is the module of global sections of the r-order jet bundle $J^rY \to X$ of $Y \to X$. Furthermore, let $Y \to X$ be an arbitrary smooth fibre bundle. An r-order (not necessarily linear) differential operator on $Y \to X$ is defined as a bundle morphism of $J^rY \to X$ to some smooth fibre bundle over X. As a consequence, Lagrangian formalism on smooth manifolds is conventionally phrased in the jet terms.

If $\mathcal{A}$ is not commutative, there are different definitions of (linear) differential operators on modules over $\mathcal{A}$ [1, 2, 3, 5, 6]. Though the notion of a jet can be extended to a module P over a noncommutative ring $\mathcal{A}$, only the first order jet module $\mathcal{J}^1(P)$ is the representative object of a certain class of first order differential operators on P.

Let $\mathcal{A}$ be a commutative $\mathcal{K}$-ring. Let P and Q be $\mathcal{A}$-modules (central bimodules). The $\mathcal{K}$-module $\mathrm{Hom}_{\mathcal{K}}(P, Q)$ of $\mathcal{K}$-linear homomorphisms $\Phi : P \to Q$ is endowed with the two different $\mathcal{A}$-module structures

$$(a\Phi)(p) := a\Phi(p), \qquad (\Phi \bullet a)(p) := \Phi(ap), \qquad a \in \mathcal{A}, \quad p \in P. \tag{2}$$

We will refer to the second one as the $\mathcal{A}^\bullet$-module structure. Let us put

$$\delta_a \Phi := a\Phi - \Phi \bullet a, \qquad a \in \mathcal{A}. \tag{3}$$

The following definitions are equivalent.

Definition 1. An element $\Delta \in \mathrm{Hom}_{\mathcal{K}}(P, Q)$ is an r-order Q-valued differential operator on P if $\delta_{a_0} \circ \cdots \circ \delta_{a_r} \Delta = 0$ for any tuple of $r+1$ elements $a_0, \ldots, a_r$ of $\mathcal{A}$.

Definition 2. An element $\Delta \in \mathrm{Hom}_{\mathcal{K}}(P, Q)$ is a zero order differential operator if $\delta_a \Delta = 0$ for all $a \in \mathcal{A}$, and Δ is a differential operator of order $r > 0$ if $\delta_a \Delta$ for all $a \in \mathcal{A}$ is an $(r-1)$-order differential operator.

Given an $\mathcal{A}$-module P, let us consider the tensor product $\mathcal{A} \otimes_{\mathcal{K}} P$ of $\mathcal{K}$-modules $\mathcal{A}$ and P. We put

$$\delta^b(a \otimes p) := (ba) \otimes p - a \otimes (bp), \qquad p \in P, \qquad a, b \in \mathcal{A}. \tag{4}$$

Let us denote by μ^{k+1} the $\mathcal{A}$-submodule of $\mathcal{A} \otimes_{\mathcal{K}} P$ generated by elements of the type

$$\delta^{b_0} \circ \cdots \circ \delta^{b_k}(\mathbf{1} \otimes p).$$

Definition 3. The k-order jet module $\mathcal{J}^k(P)$ of a module P is defined as the quotient of the $\mathcal{K}$-module $\mathcal{A} \otimes_{\mathcal{K}} P$ by μ^{k+1}. We denote its elements $a \otimes_k p$.

The $\mathcal{K}$-module $\mathcal{J}^k(P)$ is endowed with the $\mathcal{A}$- and $\mathcal{A}^\bullet$-module structures

$$b(a \otimes_k p) := ba \otimes_k p, \qquad b \bullet (a \otimes_k p) := a \otimes_k (bp). \tag{5}$$

There exists the module morphism

$$J^k : P \ni p \to \mathbf{1} \otimes_k p \in \mathcal{J}^k(P) \tag{6}$$

of the $\mathcal{A}$-module P to the $\mathcal{A}^\bullet$-module $\mathcal{J}^k(P)$ such that $\mathcal{J}^k(P)$, seen as an $\mathcal{A}$-module, is generated by elements $J^k p$, $p \in P$.

Theorem 4. Any Q-valued differential operator Δ of order k on an $\mathcal{A}$-module P factorizes uniquely

$$\Delta : P \xrightarrow{J^k} \mathcal{J}^k(P) \longrightarrow Q$$

through the morphism J^k (6) and some $\mathcal{A}$-module homomorphism $\mathfrak{f}^\Delta : \mathcal{J}^k(P) \to Q$ [4].

Let us denote $J : P \ni p \mapsto \mathbf{1} \otimes p \in \mathcal{A} \otimes P$. The proof is based on the fact that

$$\delta_{b_0} \circ \cdots \circ \delta_{b_k}(\mathfrak{f} \circ J)(p) = \mathfrak{f}(\delta^{b_0} \circ \cdots \circ \delta^{b_k}(\mathbf{1} \otimes p)) \tag{7}$$

for any $\mathfrak{f} \in \mathrm{Hom}_{\mathcal{A}}(\mathcal{A} \otimes P, Q)$. The correspondence $\Delta \mapsto \mathfrak{f}^\Delta$ yields the above mentioned isomorphism (1).

Let now a $\mathcal{K}$-ring $\mathcal{A}$ need not be commutative. Let P and Q be two-sided $\mathcal{A}$-modules. Two-sided $\mathcal{A}$-modules throughout are assumed to be central bimodules over the center $\mathcal{Z}_{\mathcal{A}}$ of $\mathcal{A}$. The $\mathcal{K}$-module $\mathrm{Hom}_{\mathcal{K}}(P, Q)$ can be provided with the left $\mathcal{A}$- and right $\mathcal{A}^\bullet$-module structures (2) and the similar right and left structures

$$(\Phi a)(p) := \Phi(p)a, \qquad (a \bullet \Phi)(p) := \Phi(pa), \qquad a \in \mathcal{A}, \qquad p \in P. \tag{8}$$

For the sake of convenience, we will refer to $\mathcal{A} - \mathcal{A}^\bullet$ structures (2) and (8) as the left and right $\mathcal{A} - \mathcal{A}^\bullet$ structures, respectively. Let us put

$$\overline{\delta}_a \Phi := \Phi a - a \bullet \Phi, \qquad a \in \mathcal{A}, \qquad \Phi \in \mathrm{Hom}_{\mathcal{K}}(P, Q). \tag{9}$$

It is readily observed that $\delta_a \circ \overline{\delta}_b = \overline{\delta}_b \circ \delta_a$ for all $a, b \in \mathcal{A}$.

We follow the notion of a differential operator in [5]. Let P and Q be regarded as left $\mathcal{A}$-modules. Let us consider the $\mathcal{K}$-module $\mathrm{Hom}_{\mathcal{K}}(P, Q)$ provided with the left $\mathcal{A} - \mathcal{A}^\bullet$ module structure (2). We denote $\mathcal{Z}_0$ its center, i.e., $\delta_a \Phi = 0$ for all $\Phi \in \mathcal{Z}_0$ and $a \in \mathcal{A}$. Let $\mathcal{I}_0 = \overline{\mathcal{Z}}_0$ be the $\mathcal{A} - \mathcal{A}^\bullet$ submodule of $\mathrm{Hom}_{\mathcal{K}}(P, Q)$ generated by $\mathcal{Z}_0$. Let us consider: (i) the quotient $\mathrm{Hom}_{\mathcal{K}}(P, Q)/\mathcal{I}_0$, (ii) its center $\mathcal{Z}_1$, (iii) the $\mathcal{A} - \mathcal{A}^\bullet$ submodule $\overline{\mathcal{Z}}_1$ of $\mathrm{Hom}_{\mathcal{K}}(P, Q)/\mathcal{I}_0$ generated by $\mathcal{Z}_1$, and (iv) the $\mathcal{A} - \mathcal{A}^\bullet$ submodule $\mathcal{I}_1$ of $\mathrm{Hom}_{\mathcal{K}}(P, Q)$ given by the relation $\mathcal{I}_1/\mathcal{I}_0 = \overline{\mathcal{Z}}_1$. Then we define the $\mathcal{A} - \mathcal{A}^\bullet$ submodules $\mathcal{I}_r$, $r = 2, \ldots,$ of $\mathrm{Hom}_{\mathcal{K}}(P, Q)$ by induction as follows:

$$\mathcal{I}_r/\mathcal{I}_{r-1} = \overline{\mathcal{Z}}_r, \tag{10}$$

where $\overline{\mathcal{Z}}_r$ is the $\mathcal{A} - \mathcal{A}^\bullet$ module generated by the center $\mathcal{Z}_r$ of the quotient $\mathrm{Hom}_{\mathcal{K}}(P, Q)/\mathcal{I}_{r-1}$.

Definition 5. Elements of the submodule $\mathcal{I}_r$ of $\mathrm{Hom}_{\mathcal{K}}(P, Q)$ are said to be the left r-order Q-valued differential operators on a two-sided $\mathcal{A}$-module P.

An equivalent definition is the following [6]

Definition 6. An element of $\Delta \in \mathrm{Hom}_{\mathcal{K}}(P, Q)$ is a left r-order Q-valued differential operator on an $\mathcal{A}$-module P if it is a finite sum

$$\Delta(p) = b_i \Phi^i(p) + \Delta_{r-1}(p), \qquad b_i \in \mathcal{A}, \tag{11}$$

where Δ_{r-1} and $\delta_a\Phi^i$ for all $a\in\mathcal{A}$ are left $(r-1)$-order differential operators if $r>0$, and they vanish if $r=0$.

The set of these operators is provided with the left and right $\mathcal{A}-\mathcal{A}^\bullet$ module structures.

If $\mathcal{A}$ is a commutative ring, Definitions 5 and 6 come to Definition 2.

If $P=Q=\mathcal{A}$, derivations of $\mathcal{A}$ and their compositions are differential operators on $\mathcal{A}$ in accordance with Definition 6 [6].

By analogy with Definitions 5 and 6, one can define right differential operators on two-sided $\mathcal{A}$-modules as follows.

Definition 7. Let P and Q be seen as right $\mathcal{A}$-modules over a noncommutative $\mathcal{K}$-ring $\mathcal{A}$. An element $\Delta\in\mathrm{Hom}_{\mathcal{K}}(P,Q)$ is said to be a right zero order Q-valued differential operator on P if it is a finite sum $\Delta=\Phi^i b_i$, $b_i\in\mathcal{A}$, where $\overline{\delta}_a\Phi^i=0$ for all $a\in\mathcal{A}$. An element $\Delta\in\mathrm{Hom}_{\mathcal{K}}(P,Q)$ is called a right differential operator of order $r>0$ on P if

$$\Delta(p)=\Phi^i(p)b_i+\Delta_{r-1}(p), \qquad b_i\in\mathcal{A}, \tag{12}$$

where Δ_{r-1} and $\overline{\delta}_a\Phi^i$ for all $a\in\mathcal{A}$ are right $(r-1)$-order differential operators.

Definition 5 and Definition 7 of differential operators on two-sided $\mathcal{A}$-modules are not equivalent, but one can combine them as follows.

Definition 8. Let P and Q be two-sided modules over a noncommutative $\mathcal{K}$-ring $\mathcal{A}$. An element $\Delta\in\mathrm{Hom}_{\mathcal{K}}(P,Q)$ is a two-sided zero order Q-valued differential operator on P if it is either a left or right zero order differential operator. An element $\Delta\in\mathrm{Hom}_{\mathcal{K}}(P,Q)$ is said to be a two-sided differential operator of order $r>0$ on P if it is brought both into the form $\Delta=b_i\Phi^i+\Delta_{r-1}$, $b_i\in\mathcal{A}$, and $\Delta=\overline{\Phi}^i\overline{b}_i+\overline{\Delta}_{r-1}$, $\overline{b}_i\in\mathcal{A}$, where Δ_{r-1}, $\overline{\Delta}_{r-1}$ and $\delta_a\Phi^i$, $\overline{\delta}_a\overline{\Phi}^i$ for all $a\in\mathcal{A}$ are two-sided $(r-1)$-order differential operators.

One can think of this definition as a generalization of the definition of first order differential operators in noncommutative geometry in [2].

It is readily observed that two-sided differential operators in Definition 8 need not be left or right differential operators, and *vice versa*. At the same time, $\mathcal{A}$-valued derivations of a $\mathcal{K}$-ring $\mathcal{A}$ and their compositions obey Definition 8.

Turn now to the notion of jets in noncommutative geometry. One can follow Definition 3 in order to define the left jets of a two-sided module P over a noncommutative ring $\mathcal{A}$. However, the relation (7) does not hold if $k>0$. Therefore, no module of left jets is the representative object of left differential operators. The notion of a right jet of P meets the similar problem.

Given an $\mathcal{A}$-module P, let us consider the tensor product $\mathcal{A}\otimes_{\mathcal{K}} P\otimes_{\mathcal{K}}\mathcal{A}$ of $\mathcal{K}$-modules $\mathcal{A}$ and P. We put

$$\begin{aligned}&\delta^b(a\otimes p\otimes c):=(ba)\otimes p\otimes c-a\otimes(bp)\otimes c,\\&\overline{\delta}^b(a\otimes p\otimes c):=a\otimes p\otimes(cb)-a\otimes(pb)\otimes c,\qquad p\in P,\qquad a,b,c\in\mathcal{A}.\end{aligned}$$

Let us denote by μ^1 the two-sided $\mathcal{A}$-submodule of $\mathcal{A}\otimes_{\mathcal{K}} P\otimes_{\mathcal{K}}\mathcal{A}$ generated by elements of the type $\overline{\delta}^c\circ\delta^b(\mathbf{1}\otimes p\otimes\mathbf{1})$. One can define the first order two-sided jet module $\mathcal{J}^1(P)=\mathcal{A}\otimes_{\mathcal{K}} P\otimes_{\mathcal{K}}\mathcal{A}/\mu^1$ of P. Let us denote

$$J:P\ni p\mapsto\mathbf{1}\otimes p\otimes\mathbf{1}\in\mathcal{A}\otimes P\otimes\mathcal{A}.$$

Then the equality

$$\delta_{b_0}\circ\cdots\circ\delta_{b_k}(\mathfrak{f}\circ J)(p)=\mathfrak{f}(\delta^{b_0}\circ\cdots\circ\delta^{b_k}(\mathbf{1}\otimes p\otimes\mathbf{1}))$$

holds for any $\mathfrak{f}\in\mathrm{Hom}_{\mathcal{A}-\mathcal{A}}(\mathcal{A}\otimes P,Q)$. It follows that $\mathcal{J}^1(P)$ is the representative object of the functor $Q\to\overline{\mathrm{Diff}}_1(P,Q)$ where the $\mathcal{K}$-module $\overline{\mathrm{Diff}}_1(P,Q)$ consists of two-sided first order differential operators Δ by Definition 8 which obey the condition $\overline{\delta}^c\circ\delta^b\Delta=0$ for all $c,b\in\mathcal{A}$. They are the first order differential operators defined in [2].

References

[1] A.Borowiec, Vector fields and differential operators: Noncommutative case, *Cech. J. Phys.* **47** (1997) 1093; *E-print arXiv*: q-alg/9710006.

[2] M.Dubois-Violette and T.Masson, On the first-order operators on bimodules, *Lett. Math. Phys.* **37** (1996) 467.

[3] P.Jara and D.Llena, Lie bracket of vector fields in noncommutative geometry, *E-print arXiv*: math.RA/0306044.

[4] I.Krasil'shchik, V.Lychagin and A.Vinogradov, *Geometry of Jet Spaces and Nonlinear Partial Differential Equations* (Gordon and Breach, Glasgow, 1985).

[5] V.Lunts and A.Rosenberg, Differential operators on noncommutative rings, *Selecta Mathematica, New Series* **3** (1997) 335.

[6] G.Sardanashvily, On the definition of higher order differential operators in noncommutative geometry, *E-print arXiv*: math-ph/0308013.